Navigating the Misconceptions: A Guide to Educating Patients on Antimicrobial resistance

Romana Abbas

University of Karachi, Faculty of Pharmacy, Karachi, Sindh, Pakistan.

ABSTRACT

Adherence to prescribed medical regimens is essential for the successful eradication of bacterial pathogens and the preservation of global health. However, in contemporary clinical practice, the premature cessation of antibiotic treatment has emerged as a critical driver of antimicrobial resistance (AMR). The phenomenon of incomplete antibiotic courses is closely linked with patient misconceptions regarding symptom relief, lack of health literacy, and the perceived burden of long-term medication. This study aims to investigate the primary socio-behavioral causes contributing to treatment of non-adherence and to analyze the subsequent biological consequences on microbial selection and resistance development. Data was collected through a structured survey exploring patient motivations for stopping medication early, awareness of AMR risks, and the self-reported frequency of leftover antibiotics. The findings highlight a significant disconnect between the clinical necessity of full-course therapy and the patient's behavior of equating "feeling better" with "being cured." By identifying the patterns and perceptions that lead to sub-lethal dosing, this research underscores the urgent need for targeted patient education and the implementation of robust stewardship programs to safeguard the efficacy of current antimicrobial therapies.

Keywords: Antimicrobial Resistance (AMR), Antibiotic Stewardship, Multi-Drug-Resistant Organisms (MDRO), Patient Compliance, Public Health Education

Received :2-2-2026 Accepted 20-04-2026 Published :21-4-2026

Email: romanaabbas055@gmail.com

INTRODUCTION

Antimicrobial resistance (AMR) is an escalating public health crisis, posing significant threats to the efficacy of treatments for infectious diseases globally. This phenomenon arises when microorganisms—such as bacteria, viruses, fungi, and parasites—develop mechanisms that render medications ineffective. The growing prevalence of AMR not only complicates treatment regimens but also results in longer hospital stays, higher medical costs, and increased mortality rates. In this article, we will explore the intricate mechanisms underlying AMR, its implications for healthcare systems, and potential strategies to combat this formidable challenge. [1-15]

Understanding AMR: A Global Health Concern

AMR is recognized as one of the most pressing issues in contemporary medicine. As antibiotics and other antimicrobials are misused or overused, pathogens evolve resistance, leading to infections that are more difficult to treat. The World Health Organization (WHO) has warned that AMR could lead to a future where minor infections become life-threatening, drastically reversing medical advancements made over the past century.

Mechanisms of Resistance: How Pathogens Adapt

Antimicrobial resistance arises through a range of adaptive mechanisms that enable microorganisms to withstand the effects of antimicrobial agents. These

mechanisms include enzymatic inactivation of antibiotics, alterations in antibiotic targets, reduced cell-wall permeability, activation of efflux pumps, and the utilization of alternative metabolic pathways. In addition to intrinsic resistance, microorganisms may acquire resistance through genetic mutations or through the transfer of resistance genes between organisms. The extensive and often inappropriate use of antimicrobial agents further intensifies selective pressure, promoting the survival and spread of resistant strains. Moreover, the ability of resistance determinants to disseminate through both vertical and horizontal gene transfer facilitates their movement across bacterial populations and even between different species. Together, these processes illustrate the dynamic and multifaceted nature of antimicrobial resistance, emphasizing the importance of responsible antimicrobial use and continued research efforts to limit its growing global impact.

Clinical Consequences of antimicrobial resistance

Antimicrobial resistance has significant clinical implications, as infections caused by resistant microorganisms are often more difficult to treat and may lead to prolonged illness, treatment failure, and increased healthcare costs. In such cases, patients may require alternative or more potent antimicrobial therapies, which can be less accessible and may carry greater risks of adverse effects. The rising prevalence of multidrug-resistant organisms further complicates treatment strategies and threatens the effectiveness of commonly used antibiotics. Consequently, antimicrobial resistance not only compromises patient outcomes but also places an increasing burden on healthcare systems, highlighting the importance of responsible antibiotic use and improved patient awareness regarding appropriate antimicrobial therapy.

Role of Antibiotic Stewardship

Antibiotic stewardship plays a critical role in limiting the emergence and spread of antimicrobial resistance by promoting the responsible and appropriate use of antimicrobial agents. Stewardship initiatives focus on ensuring that antibiotics are prescribed only when necessary, in the correct dosage, and for the appropriate duration. In addition, educating patients about the importance of completing prescribed antibiotic courses and avoiding unnecessary antibiotic use is essential to reduce behaviors that contribute to resistance development. Effective stewardship strategies therefore support both optimal patient care and the long-term preservation of antibiotic effectiveness.

Antimicrobial resistance is driven by multiple interconnected factors, including inappropriate antibiotic use, genetic adaptability of microorganisms, and the limited development of new antimicrobial agents. The increasing prevalence of resistant pathogens, particularly multidrug-resistant gram-negative organisms, poses significant challenges for clinical treatment and threatens the effectiveness of currently available therapies. Researchers emphasize that addressing AMR requires a combination of improved antimicrobial stewardship, responsible prescribing practices, and the development of innovative therapeutic strategies. Continued research and coordinated public health efforts remain essential to limit the spread of resistant organisms and preserve the efficacy of existing antimicrobial treatments.The reviewed studies indicate that antimicrobial resistance is a growing global health concern driven by the misuse and overuse of antibiotics, along with the natural ability of bacteria to evolve through mutations and horizontal gene transfer. The literature also highlights the significant clinical and economic impacts of resistant infections, including increased hospitalizations and higher healthcare costs. These findings emphasize the need for stronger antimicrobial stewardship and improved public awareness to address the continuing challenge of antimicrobial resistance.

Charani, Esmita, and Alison Holmes. "Antibiotic stewardship—Over the past two decades, significant efforts have been made to optimize antibiotic use in hospitals to address the growing threat of antimicrobial resistance. However, antibiotic decision-making often remains suboptimal due to limited resources for

stewardship programs and the influence of social and cultural factors within healthcare settings. Variations in prescribing practices across specialties and limited involvement of allied healthcare professionals further contribute to inconsistent antibiotic use. These findings highlight the need for context-specific stewardship interventions that consider professional dynamics and institutional practices to improve antibiotic use and ultimately enhance patient outcomes.

Antibiotic resistance is an escalating global threat that requires both the development of new antimicrobial agents and the more responsible use of existing antibiotics. Evidence suggests that effective mitigation strategies include regulatory measures, public health policies, and improved clinical practices aimed at promoting appropriate antibiotic use. Strengthening antimicrobial stewardship programs, increasing public awareness, and supporting healthcare professionals in evidence-based prescribing are essential steps to reduce unnecessary antibiotic use and limit the progression of antimicrobial resistance.

The reviewed literature emphasizes that multidrug-resistant organisms (MDRO) represent a critical public health threat due to their widespread prevalence, high virulence, and resistance to multiple antibiotics. Studies highlight the contribution of human and animal antibiotic use, environmental contamination, and bacterial genetic adaptability to the dissemination of MDRO. Effective strategies to combat this threat include robust surveillance systems, strict infection control measures, appropriate antibiotic therapy, and the development of novel antimicrobials and biocides. A coordinated “One Health” approach, integrating human, animal, and environmental health, is essential to limit the spread of resistant pathogens and safeguard public health.

While the discovery of new antimicrobial agents remains an important strategy, the pace of drug development lags behind the rapid evolution of multidrug-resistant bacteria. Current research emphasizes the potential of alternative approaches, including combination therapies, bacteriophage therapy, antimicrobial adjuvants, and nanotechnology, to enhance the efficacy of existing antimicrobials. These emerging strategies, alongside a clear understanding of the causes and mechanisms of AMR, offer promising avenues to curb the spread of resistance and address the growing public health threat posed by multidrug-resistant pathogens.

"Systematic literature Despite the growing prevalence of multidrug-resistant organisms (MDROs), comprehensive data on the burden and clinical outcomes of infections caused by these pathogens remain limited. The ABOUT-MDRO project seeks to address this gap by systematically evaluating the impact of key infections due to target MDROs, providing crucial insights into their epidemiology, associated morbidity and mortality, and the challenges they pose to healthcare systems. These findings highlight the urgent need for improved surveillance, targeted interventions, and strategies to mitigate the public health impact of MDRO infections.

The economic and health burden of antimicrobial resistance is substantial and projected to worsen globally. According to OECD estimates, resistant infections could cause 2.4 million deaths in Europe, North America, and Australia over the next 30 years, with annual costs reaching up to US$3.5 billion. Low- and middle-income countries are particularly vulnerable, with current resistance rates already ranging from 40–60% in nations such as Brazil, Indonesia, and Russia, and predicted to increase at a faster rate than in high-income countries. Recent European data indicate that the burden of resistant infections rivals that of influenza, tuberculosis, and HIV/AIDS combined, disproportionately affecting infants and the elderly. Alarmingly, 39% of these infections are caused by bacteria resistant to last-line antibiotics, posing serious challenges for treatment and highlighting the urgent need for effective interventions.

Comparative analyses of infections caused by antimicrobial-resistant versus susceptible bacteria indicate substantially worse outcomes for resistant infections. Across 175 reports of nosocomial and community-acquired infections, patients with drug-resistant strains experienced at least twice the mortality, hospitalization rates, and length of hospital stay compared to those with susceptible strains. These poor outcomes result from both ineffective antimicrobial therapy and complications arising from prior treatments. Collectively, these findings demonstrate that antimicrobial resistance not only poses a major public health challenge but also imposes a significant economic burden on healthcare systems.

Antimicrobial resistance (AMR) is a pressing global health challenge driven by the misuse and overuse of antibiotics in humans, animals, and agriculture, leading to the emergence and spread of resistant microorganisms. Since the discovery of penicillin, multidrug-resistant pathogens have increasingly complicated infection management, threatening routine medical procedures and posing severe risks to human and animal health. Microbial mechanisms such as enzymatic modification and biofilm formation further enhance resistance, while the economic burden of AMR is projected to reach trillions of dollars globally. Effective mitigation requires a One Health approach that integrates human, animal, and environmental perspectives, supported by robust surveillance, antimicrobial stewardship, research on new therapies, and public education. International collaboration and innovative tools, including artificial intelligence for improved diagnostics and treatment, are essential to preserve the efficacy of existing antibiotics and limit the growing impact of AMR.

While antibiotics and vaccines have historically controlled bacterial infections and improved public health, the growing threat of antimicrobial resistance underscores the need for innovative approaches. Advances in vaccine technology—including reverse vaccinology, novel adjuvants, and rationally designed bacterial outer membrane vesicles—offer opportunities to target multidrug-resistant bacteria. Complementary strategies, such as enhanced delivery of small-molecule antibacterials, modulation of host factors, inhibition of bacterial virulence, monoclonal antibodies, and microbiome interventions, also show promise. Addressing AMR effectively requires both vaccines and antibacterial therapies, supported by a coordinated research agenda and recognition of the societal value of interventions that prevent or treat life-threatening bacterial infections[16, 17].

The rise of antimicrobial resistance highlights the importance of infection prevention and control (IPC) as a critical strategy to limit the emergence and spread of resistant microorganisms. Exposure to antimicrobials allows susceptible organisms to be eliminated, leaving resistant strains to survive and propagate, which underscores the impact of antimicrobial misuse. Effective IPC measures, including hand hygiene, vaccination, and adherence to clinical protocols, can reduce the need for antibiotic use and curb the transmission of resistant infections. Healthcare professionals, particularly nurses, play a central role in promoting compliance with IPC practices, leading vaccination efforts, and ensuring the consistent implementation of preventive strategies to mitigate antimicrobial resistance.

METHOD

This study employed a quantitative descriptive research design to investigate patient perceptions, knowledge, and behaviors related to antibiotic use and antimicrobial resistance (AMR). The design was chosen to obtain measurable data on factors influencing treatment adherence, awareness of AMR, and self-reported antibiotic practices, enabling statistical analysis of trends and correlations.

Participants

The study included 270 participants from diverse demographic backgrounds, including varying ages, education levels, and healthcare experiences.

Participation was voluntary, and anonymity was maintained to ensure honest and unbiased responses.

Data Collection Tool

A structured online questionnaire was used as the primary data collection instrument. The questionnaire addressed:

Frequency and reasons for premature antibiotic discontinuation

Awareness of antimicrobial resistance and its consequences

Self-reported storage or reuse of leftover antibiotics

Sources of information regarding antibiotics and AMR

Attitudes toward completing prescribed antibiotic courses

The questions were designed to capture both behavioral patterns and subjective perceptions related to antibiotic use and AMR awareness.

Procedure

The questionnaire was distributed online via email and social media platforms. Participants were informed of the study's purpose before responding, and data collection occurred over a defined period to ensure accessibility and convenience.

Data Analysis

Responses were compiled and analyzed using descriptive statistical methods, including frequency counts and percentage distributions, to identify common behaviors and awareness levels regarding AMR. Patterns linking patient misconceptions or behaviors to incomplete antibiotic use were examined to interpret their potential impact on antimicrobial resistance.

Ethical Considerations

All participants provided informed consent before completing the survey. No personally identifiable information was collected, ensuring anonymity and adherence to ethical standards for online research

RESULT

A total of 270 responses were recorded for the survey on antibiotic use and adherence behaviors. The participants included students, working adults, and community members, representing a broad spectrum of the population exposed to antibiotics.Most respondents admitted to not always following instructions. According to figure 1nearly 70% reported stopping antibiotics before completing the full course. When feeling better, only 38% continued the full course, whereas nearly half discontinued after a few doses, and a small fraction followed advice from family or friends.




Figure 1: Survey data about antibiotic use




Figure 2:Results of survey based study showing antibiotic use

According to figure 2, Based on the survey of 270 respondents, the data highlights a significant but incomplete public understanding of antibiotic resistance. While a majority correctly identified overuse (63%) and failure to complete prescribed courses (53.7%) as primary drivers, there is a concerning lack of awareness regarding other critical factors. Specifically, fewer than one-third of participants (28.9%) recognized that using antibiotics for viral infections contributes to the problem, and only 18.1% identified the risks associated with using leftover or shared medication. These findings suggest that while basic concepts of "misuse" are understood, more targeted education is needed to clarify that antibiotics are ineffective against viruses and that self-medication remains a high-risk behavior.

These behaviors—premature discontinuation and improper use of leftover medications—allow surviving bacteria, often the most resistant, to multiply. Over time, this promotes antimicrobial resistance (AMR), making infections harder to treat and increasing the risk of treatment failure and spread of resistant strains. Continuous incomplete or improper use of antibiotics, even among people aware of its risks, thus directly contributes to long-term public health challenges[18, 19].

Professional guidance influenced adherence: According figure 3 over half of the participants indicated they would complete the course if a doctor or pharmacist explained the risks of incomplete use. Most respondents (73%) recognized that finishing an antibiotic course is very important to prevent resistance, yet behaviors often did not match this knowledge[17].

The survey highlights a gap between knowledge and practice in antibiotic use. While most participants understand the importance of completing courses, many discontinue treatment early, mainly because they feel better. These behaviors contribute directly to the development of antimicrobial resistance, underscoring the need for patient education, proper professional guidance, and public awareness campaigns.

DISCUSSION

The survey shows that stopping antibiotics early is largely driven by **misconceptions**, such as equating symptom relief with being cured, as well as practical factors like forgetfulness and side effects. Improper use of leftover antibiotics further amplifies the problem. Biologically, these behaviors create selective pressure that allows the most resistant bacteria to survive, multiplying over time. This accelerates the development of antimicrobial resistance, leading to infections that are harder to treat, longer illness duration, and increased risk of complications. Even when awareness exists, behavioral habits often override knowledge, highlighting the challenge of translating understanding into action[16, 20].

CONCLUSION

Premature discontinuation and misuse of antibiotics contribute directly to the rise of antimicrobial resistance. Long-term consequences include more difficult-to-treat infections, higher healthcare costs, and increased public health risk. Combating this issue requires not only raising awareness but also addressing behavioral drivers, emphasizing the real-world consequences of incomplete or improper antibiotic use.

REFERENCES

1. Lebentrau, S., et al., *Impact of the medical specialty on knowledge regarding multidrug-resistant organisms and strategies toward antimicrobial stewardship.* International Urology and Nephrology, 2017. **49**(8): p. 1311-1318.
2. Wong, V.W.Y., et al., *Approaches to multidrug-resistant organism prevention and control in long-term care facilities for older people: a systematic review and meta-analysis.* Antimicrobial Resistance & Infection Control, 2022. **11**(1): p. 7.
3. Wp, S.E., M. Norhidayah, and M.N.A. Ar, *Factors associated with multidrug-resistant organism (MDRO) mortality: an analysis from the national surveillance of multidrug-resistant organism, 2018-2022.* BMC Infectious Diseases, 2025. **25**(1): p. 60.
4. Majumder, M.A.A., et al., *Antimicrobial stewardship: fighting antimicrobial resistance and protecting global public health.* Infection and drug resistance, 2020: p. 4713-4738.
5. Barber, M.F. and J.R. Fitzgerald, *Mechanisms of host adaptation by bacterial pathogens.* FEMS microbiology reviews, 2024. **48**(4): p. fuae019.
6. Anderson, J.B., *Evolution of antifungal-drug resistance: mechanisms and pathogen fitness.* Nature Reviews Microbiology, 2005. **3**(7): p. 547-556.
7. Kaye, K.S., et al., *Pathogens resistant to antimicrobial agents: epidemiology, molecular mechanisms, and clinical management.* Infectious Disease Clinics, 2004. **18**(3): p. 467-511.
8. Rice, L.B., *The clinical consequences of antimicrobial resistance.* Current opinion in microbiology, 2009. **12**(5): p. 476-481.
9. Paladino, J.A., et al., *Economic consequences of antimicrobial resistance.* Surgical infections, 2002. **3**(3): p. 259-267.
10. Michael, C.A., D. Dominey-Howes, and M. Labbate, *The antimicrobial resistance crisis: causes, consequences, and management.* Frontiers in public health, 2014. **2**: p. 145.
11. MacGowan, A.P., *Clinical implications of antimicrobial resistance for therapy.* Journal of antimicrobial chemotherapy, 2008. **62**(suppl_2): p. ii105-ii114.
12. Ha, D.R., N.M. Haste, and D.P. Gluckstein, *The role of antibiotic stewardship in promoting appropriate antibiotic use.* American journal of lifestyle medicine, 2019. **13**(4): p. 376-383.
13. Manning, M.L., J. Pfeiffer, and E.L. Larson, *Combating antibiotic resistance: the role of nursing in antibiotic stewardship.* American Journal of Infection Control, 2016. **44**(12): p. 1454-1457.
14. Mendelson, M., *Role of antibiotic stewardship in extending the age of modern medicine: continuing medical education-review.* South African Medical Journal, 2015. **105**(5): p. 414-418.
15. Çakmakçi, M., *Antibiotic stewardship programmes and the surgeon's role.* Journal of Hospital Infection, 2015. **89**(4): p. 264-266.
16. Jamil, H., et al., *KNOWLEDGE, ATTITUDES, AND PRACTICES TOWARDS MULTIDRUG-RESISTANT TYPHOID IN PEDIATRIC POPULATION IN PAKISTAN.* Insights-Journal of Life and Social Sciences, 2025. **3**(6): p. 7-16.
17. Sadia, H., et al., *To prescribe or not to prescribe in the elderly: a qualitative exploration of prescribing dilemmas among Pakistani healthcare providers.* BMJ open, 2025. **15**(4): p. e094545.
18. Sattar, A., et al., *Trustworthiness of Web-Based Pharmacy Apps in Pakistan Based on the Mobile App Rating Scale: Content Analysis and Quality Evaluation.* JMIR mHealth and uHealth, 2025. **13**(1): p. e59884.
19. Sadia, H., et al., *Enhancing medication appropriateness: Insights from the STOPP (Screening Tool of Older Persons' Prescriptions) criteria version 3 on prescribing practices among the older adults in Pakistan.* Frontiers in Pharmacology, 2025. **16**: p. 1551819.
20. Naveed, S., et al., *A DEEP INSIGHT INTO INCIDENCE, PREVALENCE, COMPLETE EPIDEMIOLOGY, AND PREDICTIVE ASSOCIATIONS AMONG MULTIPLE FACTORS OF BREAST CANCER IN KARACHI, PAKISTAN: A CROSS-SECTIONAL SURVEY OF 2023.* Journal of Medical & Health Sciences Review, 2025. **2**(4).