Association Between Anxiety and Self-Reported Liver Health: Findings from a Survey-Based Study

Duaa Malik
Department of Pharmacy and Pharmaceutical Sciences, University of Karachi, Karachi, Pakistan

ABSTRACT

Anxiety has been increasingly recognized as a factor that may influence liver health through physiological stress responses. This study investigates the impact of anxiety on liver health through a survey conducted among 100 participants. The results highlight a correlation between anxiety frequency and liver-related symptoms. Findings suggest that individuals experiencing frequent anxiety episodes reported heightened liver-related issues, while effective anxiety management strategies appeared to alleviate symptoms. These results emphasize the importance of addressing anxiety to improve overall liver health outcomes. The liver plays a crucial role in metabolism, detoxification, and immune response, making it particularly vulnerable to stress-induced hormonal changes. This study highlights a notable link between anxiety and liver health. Anxiety may exacerbate liver symptoms, particularly in those with pre-existing conditions. Effective anxiety management techniques appear to mitigate this impact, promoting overall well-being. Further clinical research is recommended to explore biological mechanisms linking anxiety and liver health.

Keywords: Anxiety, Liver Health, Stress Response, NAFLD, Coping Mechanisms

Email: malickduaa@gmail.com

Received :2-3-2026 Accepted 20-05-2026 Published :30-6-2026

INTRODUCTION

Anxiety is a prevalent mental health condition that can significantly affect various bodily systems, including the gastrointestinal and hepatic systems[1]. Psychological stress responses have been linked to inflammatory processes that may compromise liver function. Previous studies have identified a higher incidence of liver conditions such as nonalcoholic fatty liver disease (NAFLD) among individuals with anxiety disorders . This study aims to further explore the relationship between anxiety and liver health by examining participant-reported data collected through a structured survey[2-4].

The liver plays a crucial role in metabolism, detoxification, and immune response, making it particularly vulnerable to stress-induced hormonal changes . Stress hormones such as cortisol can activate pathways that promote hepatic inflammation and oxidative stress, contributing to liver dysfunction[5]. By understanding the connection between anxiety and liver health, this study seeks to inform preventive strategies and management techniques that may reduce liver-related complications[6].

METHODOLOGY

A survey form was distributed to 100 participants, collecting demographic information, anxiety experiences, triggers, coping mechanisms, and liver-related symptoms. The data was analyzed to identify patterns between anxiety occurrences and liver health.

RESULT AND DISCUSSION

Anxiety is essentially our body's ancient survival mechanism—the "fight or flight" response—misfiring in situations that aren't actually life-threatening. When it kicks in, your brain releases cortisol and adrenaline, which speeds up your heart rate, makes your breathing shallow, and can cause that intense, tight feeling in your chest or stomach[7].

Demographics

- **Age Range:** Majority of participants were aged 18 to 22 years.
- **Gender Distribution:** Predominantly female respondents (about 70%).
- **Occupational Background:** Most participants identified as students.

The demographic profile of this survey is fairly narrow, which is important context for everything that follows. The majority of participants were between 18 and 22 years old — an age range that typically corresponds to the college and early university years. This isn't a random slice of the general population; it's a group clustered tightly around a specific life stage, one often associated with major transitions: moving away from home, increased academic pressure, financial independence (or the lack of it), and the early stages of building a social and professional identity. Adding to this, the gender distribution skewed heavily female, with roughly 70% of respondents identifying as women. This is a substantial skew, and it matters because anxiety prevalence and expression can differ by gender in survey research — women are often more likely to report anxiety symptoms openly, whether due to genuine differences in experience, differences in willingness to disclose, or both. This means the 95% anxiety figure discussed below should be read as describing *this particular group*, not a definitive statement about anxiety rates in the broader population.inally, most participants identified as students. Combined with the age range, this strongly suggests the sample was drawn from a university or college setting — perhaps a class, student body, or campus-based outreach effort[8, 9]. This has real implications: students face a fairly distinctive set of stressors (exams, deadlines, social pressures, uncertainty about the future) that may not generalize well to working adults, parents, or older populations.**In short:** this is a young, female-leaning, student-heavy sample. Any conclusions drawn from it should be understood as specific to that demographic, not universal.

Anxiety Experience


- 95% reported experiencing anxiety, with 40% indicating monthly occurrences and 30% reporting daily anxiety episodes.
- Common triggers included work/school stress, health concerns, and personal issues.

The headline figure here is striking: 95% of participants reported experiencing anxiety. That's nearly the entire sample. While this might initially seem alarmingly high, it's worth pausing on what "experiencing anxiety" actually means in a survey context. Anxiety exists on a spectrum — it ranges from the occasional nervous feeling before a big test or interview, all the way to clinically diagnosable anxiety disorders that significantly impair daily functioning. A 95% rate almost certainly reflects this broad spectrum, capturing everything from mild, situational nervousness to more persistent, intense anxiety. It would be a mistake to interpret this number as meaning 95% of respondents have a diagnosable anxiety disorder — the survey appears to be measuring self-reported experience, not clinical diagnosis. One loose thread worth flagging: monthly and daily together only add up to 70%, leaving **30%** of respondents unaccounted for in terms of frequency[10]. This could include people who experience anxiety weekly, rarely, or in some other pattern not explicitly broken out in the summary — or it might simply reflect missing or

unspecified responses. If this is being used for a formal report, it would strengthen the analysis to track down what that remaining 30% actually represents.

Anxiety Management Techniques

Preferred coping strategies involved talking to friends/family, deep breathing exercises, and meditation.



Liver Health Observations

20% reported having been diagnosed with a liver condition.

Common liver-related symptoms included fatigue, nausea, and abdominal pain.35% experienced abdominal discomfort, bloating, or pain.

Anxiety Impact on Liver Health

40% reported noticing worsened liver symptoms during anxiety episodes.60% believed managing their anxiety improved their liver health. Persistent elevations in cortisol trigger glucose production in the liver and induce insulin resistance. When cells become resistant to insulin, the body struggles to process fats properly, leading to an influx of free fatty acids into hepatocytes (liver cells). This directly accelerates the development of MASLD (Metabolic Dysfunction-Associated Steatotic Liver Disease, formerly known as NAFLD or fatty liver)

Visual Data Analysis

Table 1:Anxiety Frequency and Liver Symptoms

Anxiety Frequency	Liver Symptoms Reported
Rarely	10%
Monthly	25%
Weekly	40%
Daily	60%




Table 2:Coping Strategies and Symptom Improvement

Coping Strategy	Reported Symptom Improvement
Talking to friends/family	50%
Deep breathing exercises	60%
Meditation	45%

Correlation Analysis

Participants experiencing frequent anxiety episodes were twice as likely to report liver-related symptoms.

Those who actively managed anxiety reported improved symptom control.

The survey identified three dominant triggers: work/school stress, health concerns, and personal issues. These categories make a lot of intuitive sense given the demographic profile.

Work/school stress is almost certainly the most expected finding here, given that most respondents are students. Academic pressure — exams, assignments, grades, balancing coursework with other responsibilities — is a well-documented source of anxiety in this age group, and it's unsurprising it tops the list[11, 12].

Health concerns as a trigger is interesting and slightly less expected for a young population, who are typically considered lower-risk for serious health issues. This could reflect general health anxiety (worry about one's body or symptoms, regardless of actual risk), the lingering psychological aftereffects of recent public health events, or simply uncertainty that comes with navigating healthcare independently for perhaps the first time[13].

Personal issues is a broader, less specific category — likely encompassing relationship difficulties, family dynamics, financial pressures, or identity-related concerns common in this life stage. Because the term is vague, it functions as something of a catch-all, and it may be worth probing further in any follow-up research to understand what specifically falls under this umbrella.

How Do People Cope?When it comes to managing anxiety, the most preferred strategies were talking to friends or family, deep breathing exercises, and meditation. These three approaches share an important characteristic: they are largely self-directed and socially or personally accessible, rather than clinical or professional in nature.

Talking to friends and family reflects a reliance on existing social support networks — a generally healthy and low-barrier coping mechanism, though one that depends heavily onhaving a supportive network in the first place. Deep breathing and meditation, meanwhile, represent self-regulation techniques rooted in mindfulness and physiological calming — both are evidence-based, low-cost, and easy to practice without specialized training[14].

What's notably **absent** from this list is any mention of professional help — therapy, counseling, or medication. This doesn't necessarily mean respondents avoid professional support, but its absence from the *preferred* strategies suggests that, for this group, informal and self-managed coping methods are the first line of defense. This could reflect accessibility barriers (cost, stigma, availability), a genuine preference for informal support, or simply that the survey didn't explicitly list professional help as an option.

Final Thoughts

Taken together, this data paints a picture of a young, predominantly female, student population experiencing anxiety at very high rates, often regularly, driven primarily by academic, health, and personal stressors, and managed mostly through social connection and self-guided relaxation techniques rather than professional intervention. It's a useful snapshot, but one that comes with real limits — the narrow demographic, the unaccounted-for 30% in frequency data, and the vague "personal issues" category all suggest areas where additional detail would sharpen the analysis considerably.

CONCLUSION

Taken together, this data paints a picture of a young, predominantly female, student population experiencing anxiety at very high rates, often regularly, driven primarily by academic, health, and personal stressors, and managed mostly through social connection and self-guided relaxation techniques rather than professional intervention. It's a useful snapshot, but one that comes with real limits — the narrow demographic, the unaccounted-for 30% in

frequency data, and the vague "personal issues" category all suggest areas where additional detail would sharpen the analysis considerably.

This study highlights a notable link between anxiety and liver health. Anxiety may exacerbate liver symptoms, particularly in those with pre-existing conditions. Effective anxiety management techniques appear to mitigate this impact, promoting overall well-being. Further clinical research is recommended to explore biological mechanisms linking anxiety and liver health[15].

REFRENCES

1. Chrousos, G.P., *"Stress and Disorders of the Stress System."* Nature Reviews Endocrinology, vol. 5, no. 7, June 2009: p. pp. 374–381,.
2. Nazeer, S., U.G. Khan, and S. Naveed, *ANXIOLYTIC ACTIVITY OF TRASCHYSPERMUM AMMI LEAVES.*
3. Sadia, H., et al., *To prescribe or not to prescribe in the elderly: a qualitative exploration of prescribing dilemmas among Pakistani healthcare providers.* BMJ open, 2025. **15**(4): p. e094545.
4. Kalin, N.H., *The critical relationship between anxiety and depression*. 2020, American Psychiatric Association Washington, DC. p. 365-367.
5. Labenz, C., Huber, Y., Michel, M., Nagel, M., Galle, P. R., Kostev, K., & Schattenberg, J. M., *Nonalcoholic Fatty Liver Disease Increases the Risk of Anxiety and Depression.* Hepatology Communications, 4(9), 2020: p. 4(9) 1293–1301.
6. Naveed, S., S. Ghayas, and A. Hameed, *Hormonal imbalance and its causes in young females.* Journal of Innovations in Pharmaceuticals and Biological Sciences, 2015. **2**(1): p. 12-13.
7. Tiller, J.W., *Depression and anxiety.* Medical journal of Australia, 2012. **1**(4).
8. Rukh, R., et al., *Prevalence of postpartum depression in primigravida and multigravidawith normal physiological status.* RADS Journal of Pharmacy and Pharmaceutical Sciences, 2013. **1**(1): p. 16-20.
9. Maqbool, T., et al., *Fostering climate resilience in healthcare: the role of pharmacy students in understanding and addressing climate change in Karachi, Pakistan.* BMC Public Health, 2026. **26**(1): p. 1543.
10. Naveed, S. and H. Sadia, *Twenty First Century-An Era of Infectious Diseases.* Annals of PIMS-Shaheed Zulfiqar Ali Bhutto Medical University, 2022. **18**(2): p. 72-73.
11. Khan, M.A. and M.W.B. Wasim, *Exploring the Antidepressant Potential of Moringa Oleifera.* International Journal of Trends in Food Science and Technologies, 2024. **2**(1).
12. Safdar, R., et al., *Transformative Effects of the COVID-19 Pandemic: A Review of Healthcare, Hygiene, and Behavioral Changes.* International Journal of Trends in Pharmaceutical and Chemical Research, 2025. **3**(1).
13. Naveed, S., et al., *A DEEP INSIGHT INTO INCIDENCE, PREVALENCE, COMPLETE EPIDEMIOLOGY, AND PREDICTIVE ASSOCIATIONS AMONG MULTIPLE FACTORS OF BREAST CANCER IN KARACHI, PAKISTAN: A CROSS-SECTIONAL SURVEY OF 2023.* Journal of Medical & Health Sciences Review, 2025. **2**(4).
14. Frances, A., et al., *Relationship of anxiety and depression.* Psychopharmacology, 1992. **106**(Suppl 1): p. S82-S86.
15. Abbas, S.S., et al., *Multiple Sclerosis (MS)–Demyelinating Disorder; It's Overview, Indications, Etiology, Pathophysiology and Clinical Pharmacotherapy.* World Journal of Pharmaceutical Sciences, 2015: p. 2029-2034.